

University of Groningen

## **Draft Genome Sequences of 10 *Bacillus subtilis* Strains That Form Spores with High or Low Heat Resistance**

Berendsen, Erwin M.; Wells-Bennik, Marjon H.J.; Krawczyk, Antonina O.; de Jong, Anne; van Heel, Auke; Eijlander, Robyn T.; Kuipers, Oscar P.

*Published in:*  
Genome Announcements

*DOI:*  
[10.1128/genomeA.00124-16](https://doi.org/10.1128/genomeA.00124-16)

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2016

[Link to publication in University of Groningen/UMCG research database](#)

### *Citation for published version (APA):*

Berendsen, E. M., Wells-Bennik, M. H. J., Krawczyk, A. O., de Jong, A., van Heel, A., Eijlander, R. T., & Kuipers, O. P. (2016). Draft Genome Sequences of 10 *Bacillus subtilis* Strains That Form Spores with High or Low Heat Resistance. *Genome Announcements*, 4(2), 1-2. [e00124-16].  
<https://doi.org/10.1128/genomeA.00124-16>

### **Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

### **Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

# Draft Genome Sequences of 10 *Bacillus subtilis* Strains That Form Spores with High or Low Heat Resistance

Erwin M. Berendsen,<sup>a,b,c</sup> Marjon H. J. Wells-Bennik,<sup>b,c</sup> Antonina O. Krawczyk,<sup>a,c</sup> Anne de Jong,<sup>a,c</sup> Auke van Heel,<sup>a</sup> Robyn T. Eijlander,<sup>a,c</sup> Oscar P. Kuipers<sup>a,c</sup>

Molecular Genetics, University of Groningen, Groningen, Groningen, The Netherlands<sup>a</sup>; NIZO food research, Ede, The Netherlands<sup>b</sup>; Top Institute Food and Nutrition (TIFN), Wageningen, The Netherlands<sup>c</sup>

**Here, we report the draft genome sequences of 10 isolates of *Bacillus subtilis*, a spore forming Gram-positive bacterium. The strains were selected from food products and produced spores with either high or low heat resistance.**

Received 29 January 2016 Accepted 6 February 2016 Published 17 March 2016

**Citation** Berendsen EM, Wells-Bennik MHJ, Krawczyk AO, de Jong A, van Heel A, Eijlander RT, Kuipers OP. 2016. Draft genome sequences of 10 *Bacillus subtilis* strains that form spores with high or low heat resistance. *Genome Announc* 4(2):e00124-16. doi:10.1128/genomeA.00124-16.

**Copyright** © 2016 Berendsen et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Oscar P. Kuipers, o.p.kuipers@rug.nl.

*Bacillus subtilis* is a soil-dwelling organism that can also be found in the mammalian microbiota (1). *B. subtilis* strains can be used as cell factories for enzyme or metabolite production, as biocontrol agents, or even as probiotics (2). However, they can also form low- or high-heat-resistant spores that may survive food processing techniques and cause food spoilage in consumer products. Ten strains of *B. subtilis* were isolated as spores from food products (Table 1). The spore heat resistance traits of some of these strains were described in a study by Berendsen et al. (3). The sequences of these strains will provide valuable information on genes involved in sporulation and germination (4, 5). Moreover, these and previously published *B. subtilis* strains can be grouped according to phenotype, and subsequent gene trait matching can be used to identify genes involved in the phenotype of interest.

The 10 strains were grown overnight in 10 ml of brain heart infusion (BHI) broth (Difco) at 37°C and harvested at the exponential-growth phase. After centrifugation, the cell pellet was resuspended in SET buffer (75 mM NaCl, 25 mM EDTA, 20 mM Tris-HCl [pH 7.5]) and incubated with lysozyme (2 mg/ml) and RNase (0.4 mg/ml) for 30 min at 37°C. Subsequently, the sample was treated with SDS (1% final concentration) and proteinase K (0.5 mg/ml) at 55°C for 60 min. Genomic DNA was extracted from the lysate with phenol-chloroform, precipitated with isopro-

panol and sodium acetate (300 mM), and dissolved in Tris-EDTA (TE) buffer. The isolated DNA was sheared to 500-bp fragments in the Covaris (KBiosciences) ultrasonic device for preparing the next-generation sequencing (NGS) library preps using the paired-end NEB NextGen library preparation kit. The libraries were 101-base paired-end sequenced on an Illumina HiSeq 2000 by multiplexing 12 samples per flow cell. Velvet (6) was used to perform a *de novo* paired-end assembly on each of the 10 genomes, resulting in the draft genome sequences (Table 1). Annotation of the genomes was done using the following steps: (i) scaffolds were uploaded to the RAST server (7) and automatically annotated using the SEED bases method on this server, (ii) the resulting annotated scaffolds were mapped using CONTIGuator (8) on their closest neighbor (identified by RAST) to generate the pseudogenome, (iii) locus tags were added to each feature using an in-house-developed Perl script, according to the NCBI standard, (iv) BAGEL3 (9) was used to find and annotate bacteriocin gene clusters, and (v) the protein annotation was extended using InterProScan (10).

**Nucleotide sequence accession numbers.** The genome sequence of the 10 *B. subtilis* strains have been deposited as whole-genome shotgun projects at DDBJ/EMBL/GenBank under the accession numbers listed in Table 1.

TABLE 1 Genome features and GenBank accession numbers of the strains

Strain ID	Strain	Isolate source	Accession no.	Genome size (Mb)	Coverage (×)
B4067	<i>B. subtilis</i> A163	Chicken soup	<a href="https://genbank.ncbi.nlm.nih.gov/GenBank/FASTA/JXHS0000000000.fna">JXHS00000000</a>	4.31	1,168
B4068	<i>B. subtilis</i> CC2	Curry cream	<a href="https://genbank.ncbi.nlm.nih.gov/GenBank/FASTA/JXHK0000000000.fna">JXHK00000000</a>	3.98	227
B4069	<i>B. subtilis</i> IIC14	Binding flour	<a href="https://genbank.ncbi.nlm.nih.gov/GenBank/FASTA/JXHL0000000000.fna">JXHL00000000</a>	4.09	434
B4070	<i>B. subtilis</i> A162	Peanut chicken soup	<a href="https://genbank.ncbi.nlm.nih.gov/GenBank/FASTA/JXHM0000000000.fna">JXHM00000000</a>	4.28	237
B4071	<i>B. subtilis</i> CC16	Curry cream soup	<a href="https://genbank.ncbi.nlm.nih.gov/GenBank/FASTA/JXHN0000000000.fna">JXHN00000000</a>	4.2	323
B4072	<i>B. subtilis</i> RL45	Red lasagna sauce	<a href="https://genbank.ncbi.nlm.nih.gov/GenBank/FASTA/JXHO0000000000.fna">JXHO00000000</a>	4.09	197
B4073	<i>B. subtilis</i> MC85	Curry soup	<a href="https://genbank.ncbi.nlm.nih.gov/GenBank/FASTA/JXHP0000000000.fna">JXHP00000000</a>	4.13	285
B4143	<i>B. subtilis</i>	Surimi	<a href="https://genbank.ncbi.nlm.nih.gov/GenBank/FASTA/JXLQ0000000000.fna">JXLQ00000000</a>	4.2	311
B4145	<i>B. subtilis</i>	Cereals	<a href="https://genbank.ncbi.nlm.nih.gov/GenBank/FASTA/JXHQ0000000000.fna">JXHQ00000000</a>	4.4	258
B4146	<i>B. subtilis</i>	Mayonnaise	<a href="https://genbank.ncbi.nlm.nih.gov/GenBank/FASTA/JXHR0000000000.fna">JXHR00000000</a>	4.26	313

## ACKNOWLEDGMENTS

We thank the NGS sequence facility of the University Medical Center of Groningen (UMCG) for performing the sequencing of the strains. We also thank TIFN for contributing to the funding of the project.

## REFERENCES

1. Kunst F, Ogasawara N, Moszer I, Albertini AM, Alloni G, Azevedo V, Bertero MG, Bessi eres P, Bolotin A, Borchert S, Borriss R, Boursier L, Brans A, Braun M, Brignell SC, Bron S, Brouillet S, Bruschi CV, Caldwell B, Capuano V, Carter NM, Choi SK, Cordani JJ, Connerton IF, Cummings NJ, Daniel RA, Denzot F, Devine KM, D usterh oft A, Ehrlich SD, Emmerson PT, Entian KD, Errington J, Fabret C, Ferrari E, Foulger D, Fritz C, Fujita M, Fujita Y, Fuma S, Galizzi A, Galleron N, Ghim SY, Glaser P, Goffeau A, Golightly EJ, Grandi G, Guiseppi G, Guy BJ, et al. 1997. The complete genome sequence of the Gram-positive bacterium *Bacillus subtilis*. *Nature* 390:249–256. <http://dx.doi.org/10.1038/36786>.
2. Mongkolthanaruk W. 2012. Classification of *Bacillus* beneficial substances related to plants, humans and animals. *J Microbiol Biotechnol* 22:1597–1604. <http://dx.doi.org/10.4014/jmb.1204.04013>.
3. Berendsen EM, Zwietering MH, Kuipers OP, Wells-Bennik MHJ. 2015. Two distinct groups within the *Bacillus subtilis* group display significantly different spore heat resistance properties. *Food Microbiol* 45:18–25. <http://dx.doi.org/10.1016/j.fm.2014.04.009>.
4. Luu S, Setlow P. 2014. Analysis of the loss in heat and acid resistance during germination of spores of *Bacillus* species. *J Bacteriol* 196:1733–1740. <http://dx.doi.org/10.1128/JB.01555-14>.
5. Eijlander RT, de Jong A, Krawczyk AO, Holsappel S, Kuipers OP. 2014. SporeWeb: an interactive journey through the complete sporulation cycle of *Bacillus subtilis*. *Nucleic Acids Res* 42:D685–D691. <http://dx.doi.org/10.1093/nar/gkt1007>.
6. Zerbino DR, Birney E. 2008. Velvet: algorithms for *de novo* short read assembly using de Bruijn graphs. *Genome Res* 18:821–829. <http://dx.doi.org/10.1101/gr.074492.107>.
7. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: Rapid Annotations using Subsystems Technology. *BMC Genomics* 9:75. <http://dx.doi.org/10.1186/1471-2164-9-75>.
8. Galardini M, Biondi EG, Bazzicalupo M, Mengoni A. 2011. CONTIGuator: a bacterial genomes finishing tool for structural insights on draft genomes. *Source Code Biol Med* 6:11. <http://dx.doi.org/10.1186/1751-0473-6-11>.
9. van Heel AJ, de Jong A, Montalb n-L pez M, Kok J, Kuipers OP. 2013. BAGEL3: automated identification of genes encoding bacteriocins and (non-)bactericidal posttranslationally modified peptides. *Nucleic Acids Res* 41:W448–W453. <http://dx.doi.org/10.1093/nar/gkt391>.
10. Jones P, Binns D, Chang HY, Fraser M, Li W, McAnulla C, McWilliam H, Maslen J, Mitchell A, Nuka G, Pesseat S, Quinn AF, Sangrador-Vegas A, Scheremetjew M, Yong SY, Lopez R, Hunter S. 2014. InterProScan 5: genome-scale protein function classification. *Bioinformatics* 30:1236–1240.